

Data presented as: Percent Susceptible (# of Isolates Tested)	Penicillins			Cephalosporins		Tetracyclines	Quinolones		Other							# of all isolates tested (# of hospitals reporting)
	Ampicillin	Oxacillin	Penicillin	Ceftriaxone	Ceftaroline	Doxycycline	Ciprofloxacin	Levofloxacin	Clindamycin	Daptomycin	Erythromycin	Linezolid	Nitrofurantoin	Trimethoprim/ Sulfamethoxazole	Vancomycin	
<i>Enterococcus</i> spp.	90 17,867	R	71 4,189	R	R	46 864	57 3,671	71 3,035	R	89 2,193	19 3,040	99 16,405	87 13,404	R	90 18,039	18,553 51
<i>Enterococcus faecalis</i>	99 15,722	R	98 11,373	R	R	27 1,900	69 8,821	73 9,940	R	95 4,211	9 5,828	97 13,492	98 12,116	R	94 17,422	17,307 80
<i>Enterococcus faecium</i>	17 4,120	R	15 3,042	R	R	49 481	11 2,290	15 2,279	R	79 1,540	4 1,520	96 3,963	27 2,657	R	34 4,485	4,453 72
<i>Staphylococcus aureus</i> (all)	2 6,952	63 45,427	10 15,858	-	100 3,044	96 5,830	54 19,544	52 16,734	71 47,730	98 9,054	44 24,209	100 44,968	99 16,073	95 48,560	100 50,031	46,679 71
Methicillin-resistant <i>Staphylococcus aureus</i> (MRSA)	R	0 15,418	R	R	98 857	90 2,360	13 8,839	13 7,959	61 17,738	97 4,070	14 9,926	98 16,334	99 5,076	91 18,372	100 18,480	18,465 68
Methicillin-susceptible <i>Staphylococcus aureus</i> (MSSA)	3 3,849	99 25,794	18 8,768	-	100 2,211	97 4,354	84 10,998	85 8,843	78 28,656	99 4,321	65 12,899	99 26,807	99 7,703	97 29,084	100 29,370	29,545 61
<i>Streptococcus agalactiae</i> (Group B Strep)	84 1,207	-	100 7,521	99 557	100 86	-	-	94 845	49 7,529	100 40	46 614	100 836	-	-	100 7,865	8,290 53
<i>Streptococcus pyogenes</i> (Group A Strep)	100 412	-	99 395	100 245	-	-	-	97 164	84 442	-	87 453	100 151	-	-	100 465	479 32
<i>Streptococcus pneumoniae</i> (non-meningitis)	92 36	-	94 694	96 687	-	94 32	-	95 490	80 478	-	66 581	98 173	-	78 576	99 755	818 48
<i>Streptococcus pneumoniae</i> (meningitis)	-	-	70 404	86 455	-	-	-	-	-	-	61 84	-	-	-	-	492 29