

***Enterococcus* spp.**  
**(n=18,553 from 51 Hospitals)**

	<b>% Susceptible (IQR)</b>	<b>Number of isolates tested</b>	<b>Number of hospitals</b>
<b>Ampicillin</b>	90% (74-91)	17,867	50
<b>Penicillin</b>	71% (50-81)	4,189	29
<b>Levofloxacin</b>	71% (50-82)	3,050	29
<b>Doxycycline</b>	46% (38-48)	864	4
<b>Ciprofloxacin</b>	57% (51-79)	3,671	27
<b>Daptomycin</b>	88% (85-98)	2,193	17
<b>Erythromycin</b>	19% (10-42)	3,040	23
<b>Linezolid</b>	99% (97-100)	16,405	45
<b>Nitrofurantoin</b>	87% (71-94)	13,404	42
<b>Vancomycin</b>	90% (56-88)	18,039	51

***Enterococcus faecalis***  
(n=17,307 from 80 Hospitals)

	<b>% Susceptible (IQR)</b>	<b>Number of isolates tested</b>	<b>Number of hospitals</b>
<b>Ampicillin</b>	99% (99-100)	15,722	77
<b>Penicillin</b>	96% (96-99)	11,373	51
<b>Levofloxacin</b>	73% (69-80)	9,940	58
<b>Doxycycline</b>	27% (20-28)	1,900	10
<b>Ciprofloxacin</b>	69% (62-80)	8,821	55
<b>Daptomycin</b>	95% (95-100)	4,211	23
<b>Erythromycin</b>	9% (5-13)	5,828	43
<b>Linezolid</b>	97% (97-100)	13,492	74
<b>Nitrofurantoin</b>	98% (97-100)	12,116	71
<b>Vancomycin</b>	94% (94-98)	17,422	80

***Enterococcus faecium***  
**(n=4,453 from 72 Hospitals)**

	<b>% Susceptible (IQR)</b>	<b>Number of isolates tested</b>	<b>Number of hospitals</b>
<b>Ampicillin</b>	17% (10-27)	4,120	67
<b>Penicillin</b>	15% (7-27)	3,042	48
<b>Levofloxacin</b>	15% (7-20)	2,279	55
<b>Doxycycline</b>	49% (48-57)	481	10
<b>Ciprofloxacin</b>	11% (5-16)	2,290	52
<b>Daptomycin</b>	79% (39-97)	1,540	22
<b>Erythromycin</b>	4% (0-5)	1,520	37
<b>Linezolid</b>	96% (97-100)	3,936	67
<b>Nitrofurantoin</b>	27% (15-40)	2,657	68
<b>Vancomycin</b>	34% (27-50)	4,485	72

***Staphylococcus aureus* (ALL)**  
**(n=49,679 from 71 Hospitals)**

	<b>% Susceptible (IQR)</b>	<b>Number of isolates tested</b>	<b>Number of hospitals</b>
<b>Oxacillin</b>	63% (46-64)	45,427	61
<b>Penicillin</b>	10% (0-19)	15,858	39
<b>Doxycycline</b>	96% (98-100)	5,830	11
<b>Ceftaroline</b>	100% (100-100)	3,044	5
<b>Ciprofloxacin</b>	54% (41-61)	19,544	44
<b>Clindamycin</b>	71% (61-72)	47,730	70
<b>Daptomycin</b>	98% (100-100)	9,054	27
<b>Erythromycin</b>	44% (38-48)	24,209	54
<b>Levofloxacin</b>	52% (45-66)	16,734	47
<b>Linezolid</b>	100% (100-100)	44,968	62
<b>Nitrofurantoin</b>	99% (99-100)	16,073	57
<b>Trimethoprim/Sulfamethoxazole</b>	95% (93-97)	48,560	70
<b>Vancomycin</b>	100% (100-100)	50,031	71

**Methicillin-Resistant *Staphylococcus aureus* (MRSA)**  
(n=18,465 from 68 Hospitals)

	<b>% Susceptible (IQR)</b>	<b>Number of isolates tested</b>	<b>Number of hospitals</b>
<b>Oxacillin</b>	0%	15,418	53
<b>Levofloxacin</b>	13% (10-16)	7,959	46
<b>Ceftaroline</b>	98% (99-100)	857	5
<b>Ciprofloxacin</b>	13% (10-16)	8,839	45
<b>Clindamycin</b>	61% (48-61)	17,738	67
<b>Daptomycin</b>	97% (99-100)	4,070	27
<b>Doxycycline</b>	90% (86-100)	2,360	12
<b>Erythromycin</b>	14% (12-17)	9,926	55
<b>Linezolid</b>	98% (100-100)	16,464	55
<b>Nitrofurantoin</b>	99% (99-100)	5,076	60
<b>Trimethoprim/Sulfamethoxazole</b>	91% (89-95)	18,372	67
<b>Vancomycin</b>	100% (100-100)	18,480	68

**Methicillin-Susceptible *Staphylococcus aureus* (MSSA)**  
(n=29,545 from 61 Hospitals)

	<b>% Susceptible (IQR)</b>	<b>Number of isolates tested</b>	<b>Number of hospitals</b>
<b>Oxacillin</b>	99% (100-100)	25,794	48
<b>Penicillin</b>	18% (0-27)	8,768	34
<b>Levofloxacin</b>	85% (80-90)	8,843	45
<b>Doxycycline</b>	97% (92-100)	4,354	12
<b>Ceftaroline</b>	100% (100-100)	2,211	5
<b>Ciprofloxacin</b>	84% (81-88)	10,998	40
<b>Clindamycin</b>	78% (72-80)	28,656	61
<b>Daptomycin</b>	99% (99-100)	4,321	17
<b>Erythromycin</b>	65% (61-69)	12,899	51
<b>Linezolid</b>	99% (100-100)	26,807	51
<b>Nitrofurantoin</b>	99% (100-100)	7,703	55
<b>Trimethoprim/Sulfamethoxazole</b>	97% (96-99)	29,084	59
<b>Vancomycin</b>	100% (100-100)	29,370	61

***Streptococcus agalactiae* (Group B *Streptococcus*)**  
**(n=8,290 from 53 Hospitals)**

	<b>% Susceptible (IQR)</b>	<b>Number of isolates tested</b>	<b>Number of hospitals</b>
<b>Penicillin</b>	100% (100-100)	7,521	38
<b>Ceftriaxone</b>	99% (100-100)	557	26
<b>Ceftaroline</b>	100% (100-100)	86	1
<b>Levofloxacin</b>	94% (92-100)	845	37
<b>Clindamycin</b>	49% (7-51)	7,529	44
<b>Daptomycin</b>	100% (100-100)	40	2
<b>Erythromycin</b>	46% (24-52)	614	29
<b>Linezolid</b>	100% (100-100)	836	28
<b>Vancomycin</b>	100% (100-100)	7,865	53

***Streptococcus pyogenes* (Group A *Streptococcus*)**  
**(n=479 from 32 Hospitals)**

	<b>% Susceptible (IQR)</b>	<b>Number of isolates tested</b>	<b>Number of hospitals</b>
<b>Ampicillin</b>	100% (100-100)	412	24
<b>Penicillin</b>	99% (100-100)	395	24
<b>Ceftriaxone</b>	100% (100-100)	245	26
<b>Levofloxacin</b>	97% (100-100)	164	24
<b>Clindamycin</b>	84% (86-100)	442	27
<b>Erythromycin</b>	87% (80-100)	453	30
<b>Linezolid</b>	100% (100-100)	151	19
<b>Vancomycin</b>	100% (100-100)	465	32



***Streptococcus pneumoniae***  
**(n=1,310 from 48 Hospitals)**

	<b>% Susceptible (IQR)</b>	<b>Number of Isolates tested</b>	<b>Number of Hospitals</b>
<b>Ampicillin</b>	92% (89-100)	36	4
<b>Penicillin</b>			
Non-meningitis	94% (96-100)	694	45
Meningitis	70% (44-88)	404	27
<b>Ceftriaxone</b>			
Non-meningitis	96% (100-100)	687	44
Meningitis	86% (75-100)	455	28
<b>Levofloxacin</b>	95% (95-100)	490	36
<b>Doxycycline</b>	94% (93-98)	32	3
<b>Clindamycin</b>	80% (73-100)	478	36
<b>Erythromycin</b>			
Non-meningitis	66% (58-100)	581	46
Meningitis	61% (54-70)	84	8
<b>Linezolid</b>	98% (100-100)	173	22
<b>Trimethoprim/Sulfamethoxazole</b>	78% (68-100)	576	42
<b>Vancomycin</b>	99% (100-100)	755	48