Breastfeeding	Basics for	r CPSP	Providers
Post-Test			

Name:
Clinic:
Phone:
Email:
Date Training Completed:

True or False

- 1. Formula fed infants get more ear infections than breastfed infants.
- 2. It is cheaper to breast feed a baby than to buy formula.
- 3. Breastfeeding only benefits the baby during the first year of life.
- 4. Women who breast feed are more likely to get breast cancer.
- 5. The amount of milk a woman can make depends on the size of her breasts.

Please circle the correct answer

- 6. Identify the three factors that can increase breastfeeding duration:
 - a. Rooming-in, in-hospital exclusive breastfeeding, breastfeeding within the first hour after birth
 - b. Priming his sucking reflex with a pacifier, rooming-in, in-hospital exclusive breastfeeding
 - c. Rooming-in, swaddling to calm infant prior to breastfeeding, referral phone number for breastfeeding help after discharge
 - d. Rooming-in, encouraging family visitors, in-hospital exclusive breastfeeding
 - e. All of the above
- 7. The following active listening technique(s) will help to build the relationship between a pregnant or breastfeeding mother and the CPSP practitioner:
 - a. Open-ended questions
 - b. Affirmations
 - c. Close-ended questions
 - d. A & B only
 - e. All of the above
- 8. Two hormones that impact lactation are:
 - a. Adrenaline, oxytocin
 - b. Oxytocin, prolactin
 - c. Progesterone, testosterone
 - d. Insulin, gastrin
 - e. Pitocin, prolactin

- 9. Identify three indicators of an effective latch:
 - a. Cheeks dimpling in while sucking, nose deep into breast and gliding jaw movements
 - b. Chin deep into breast, wide angled mouth opening, rhythmic sucking bursts with swallows
 - c. Much of the areola taken into the mouth, lips flanged back by breast and cheeks dimpling in while sucking
 - d. Baby's mouth centered on the areola, audible smacking sounds and lips flanged back by breast
 - e. Audible smacking sounds, tongue visible under areola and evidence of milk in baby's mouth
- 10. A healthy newborn should nurse how many times in 24 hours?
 - a. Every 2-3 hours
 - b. 10 or more times
 - c. As often as the mother feels able
 - d. Whenever the baby cries
- 11. The <u>most</u> common reason mothers stop breastfeeding before they had planned or wanted to is:
 - a. Sore nipples
 - b. Mother returns to school or work
 - c. Lack of infant weight gain
 - d. Perceived low milk supply
 - e. Engorgement
- 12. The treatment for sore nipples includes all of the following EXCEPT:
 - a. Assessment and correction of the infant's latch at breast
 - b. Limiting nursing time
 - c. Avoid wearing clothes or bras that are too tight
 - d. Breast milk applied to nipples after feeding

THIS SECTION TO BE COMPLETED
BY LA COUNTY CPSP STAFF

Score: <u>/12</u> = <u>%</u>

Please submit your completed test to LA County CPSP: Email: cpsp@ph.lacounty.gov

Follow-up call/email completed: □YES