


What is Chlamydia?



Your sex partner just got treatment for chlamydia. For your health, you need to get treated too!

Chlamydia [kluh-mid-ee-uh] is a sexually transmitted disease (STD) that you can get from having vaginal, anal or oral sex. Because your partner is infected, there's a good chance that you are too. Many people with an STD do not have signs. This means you can still be infected even if you feel okay.

It's very important that you get treated right away. If chlamydia isn't treated, it can spread and cause serious damage in the body.

The good news is that chlamydia is easy to treat! **The best way to take care of yourself is to see a doctor right away.** But if you can't get to a doctor within one week of getting this brochure, you can take this medicine instead.

Treatment for Chlamydia

The treatment for chlamydia is a single dose of an antibiotic called

Azithromycin (*Zithromax*) 1g
(taken as four 250mg pills)

This medicine only treats chlamydia. So, it's important to get tested for other STDs as soon as you can. Having chlamydia can increase your risk of getting HIV, so make sure that an HIV test is included as part of your STD check-up.

The LA County STD Clinics offer free and confidential testing and treatment. For a clinic near you, call the STD Hotline at 1-800-758-0880 or visit www.publichealth.lacounty.gov/std

If you're pregnant or think you might be, these pills are safe but it's better for you to get a full check-up instead of taking this medicine. If you do take the pills, see a doctor 3 weeks after taking them to make sure you've been cured.

If you are pregnant and do not have a prenatal doctor, call 1-877-597-4777 for information about free or low cost prenatal care.

Important Warnings

It's safe for most people to take the pills without seeing a doctor.

DO NOT take this medicine if:

- You have a serious illness like kidney, heart or liver disease
- You've ever had a bad reaction after taking Azithromycin (Zithromax) or other antibiotics.
- Females: you have lower belly pain, pain during sex, throwing up, or fever
- Males: you have pain or swelling in the testicles (balls) or fever

If any of the above are true, or if you're not sure, see a doctor right away for treatment. If you don't have a doctor, go to one of the LA County STD clinics for free and confidential testing and treatment.



STOP

DO NOT take these pills if you've ever had a bad reaction, rash, breathing problems, or allergic reaction after taking the antibiotic azithromycin. People who are allergic to some antibiotics may be allergic to others. If you have allergies to antibiotics, check with your doctor before taking this medicine.

How to Take the Medicine

STOP Don't take these pills if you've ever had a bad reaction to antibiotics. See "Important Warnings."

Take all of the pills together, with water.

You must take all the pills to be cured.

- Take the pills with food to avoid an upset stomach.
- DO NOT take antacids (like Tums, Rolaids, or Maalox) 1 hour before or 2 hours after taking the pills. They can keep the medicine from working.
- DO NOT share this medicine or give it to anyone else. If you have other partner(s), tell them you're getting treated so that they can get checked and treated too!

Side Effects

These pills are very safe and rarely cause problems. If problems do happen, they may include rash, feeling tired, getting a yeast infection, sunburning easily, upset stomach, throwing up*, and/or diarrhea. These effects are usually not severe and don't last long.

*If you throw up within 2 hours of taking this medicine, it won't work. Go to a doctor or clinic to get more.

Allergic reactions

Although rare, a severe allergic reaction can include: a hard time breathing/tightness in the chest; closing of your throat; swelling of your face, lips or tongue; and/or hives (itchy bumps or welts on your skin). If you have any of these symptoms, call 911 or go to the nearest emergency room right away. After you get treated, call the clinic or the STD Hotline to report your reaction.

Next Steps

- **Don't have sex until 7 days after you and your partner(s) have taken the medicine** because it takes 7 days for chlamydia to go away. If you have oral, anal or vaginal sex during this time without a condom, or if the condom breaks, you can pass the infection to your partners.
- **See a doctor for a full STD check-up** as soon as you can.
- **If you have questions about the medicine**, call the clinic or doctor listed on your prescription or call the STD Hotline.
- **If you have other sex partners**, tell them you were exposed to chlamydia.
- **For more information on STDs and tips on talking to partners**, go to www.ashastd.org and www.inspotla.org

For information, free or low-cost testing, and free condoms, call the STD Hotline at:
1-800-758-0880
or visit
www.publichealth.lacounty.gov/std

Partner

Treatment



Medicine to Treat Chlamydia

IMPORTANT!
Please read ALL of these instructions carefully!

